

Equivalence of dynamical systems by bisimulation*

Arjan van der Schaft[†]

Department of Applied Mathematics, University of Twente

P.O. Box 217, 7500 AE Enschede, The Netherlands

Phone +31-53-4893449, Fax +31-53-4893800

E-mail a.j.vanderschaft@math.utwente.nl

Abstract

A general notion of bisimulation is defined for linear input-state-output systems, using analogies with the theory of concurrent processes. A characterization of bisimulation and an algorithm for computing the maximal bisimulation relation is derived using geometric control theory. Bisimulation is shown to be a notion which unifies the concepts of state space equivalence and state space reduction, and which allows to study equivalence of systems with non-minimal state space dimension. The notion of bisimulation is especially powerful for 'non-deterministic' dynamical systems, and leads in this case to a notion of equivalence which is finer than equality of external behavior. For abstractions of systems it is shown how the results specialize to previously obtained results by other authors. Extensions of the main results to the nonlinear case are provided.

Keywords State space equivalence, bisimulation, controlled invariance, non-minimal systems, maximal bisimulation relation, non-deterministic systems, abstraction, nonlinear systems.

1 Introduction

A common theme in theoretical computer science (in particular, the theory of distributed processes and computer-aided verification) and in systems and control theory is to characterize systems which are 'externally equivalent'. The intuitive idea is that we only want to distinguish between two systems if the distinction can be detected by an external system interacting with these systems. This is a fundamental notion in design, enabling us to take a 'divide and rule' strategy, and in analysis, allowing us to switch between externally equivalent representations of the same system and to reduce sub-systems to externally equivalent but simpler sub-systems.

More specifically, a main issue in the theory of distributed processes and computer-aided verification is to develop a mathematical framework that can handle the state explosion in complex systems. A crucial notion in this endeavor is the concept of *bisimulation* which

*To appear in IEEE Transactions on Automatic Control, December 2004

[†]Research partially supported by the EU-project HYBRIDGE, IST-2001-32460

expresses when a sub-process can be considered to be externally equivalent to another (hopefully simpler) process. On the other hand, classical notions in systems and control theory are *state space equivalence* of dynamical systems, and *reduction* of a dynamical system to an equivalent system with minimal state space dimension. These notions have been instrumental in e.g. linking input-output models to state space models, and in studying the properties of interconnected systems.

Developments in both areas have been rather independent ('concurrently but without communication'), one of the reasons being that the mathematical formalisms for describing both types of systems (discrete processes on the one hand, and continuous dynamical systems on the other hand) are rather different. However, with the rise of interest in *hybrid systems*, which are systems with interacting discrete and continuous dynamics, and with the renewed interest in complexity in systems theory, this situation seems not very productive, as has been argued before by several authors.

The aim of this paper is to make another step in this reapproachment between the theory of concurrent processes and mathematical systems theory by defining and studying in detail a notion of *bisimulation for continuous dynamical systems*, and to relate it to the common system-theoretic notions of state space equivalence and state space reduction.

Such an extension of the notion of bisimulation to continuous dynamical systems has been explored before in a series of innovative papers by Pappas and co-authors [13, 14, 2, 18, 19, 20, 9, 29], and the present paper is very much inspired by this work. The main difference is that in [18, 19, 20, 29] the focus is on characterizing bisimulation of a dynamical system by a 'projected' dynamical system with lower state space dimension ('an abstraction'; see Section 5), while in the present paper we deal with a general notion of bisimulation between two continuous dynamical systems and we elaborate on the relations with the system-theoretic notions of state space equivalence and state space reduction. (We note that a general definition of bisimulation relations for dynamical systems has been given before in [9].)

In the main part of this paper we study continuous dynamical systems of the form

$$\begin{aligned} \dot{x} &= Ax + Bu + Gd, & x \in \mathbb{R}^n, u \in \mathbb{R}^m, d \in \mathbb{R}^s \\ y &= Cx, & y \in \mathbb{R}^p \end{aligned} \tag{1}$$

where x are as usual the state variables, u are the input variables, y are the output variables, while d are additional input variables, which can be thought of as *disturbances*. The basic problem we address is when two systems of the form (1) can be considered to be externally *equivalent*, in the sense that for all time instants t_0 the solution trajectories for $t \geq t_0$ of one system are mimicked by the other in such a way that the input and output trajectories $u(t)$ and $y(t)$ of both systems are the same for $t \geq t_0$, *without* imposing any relation between the values of their disturbance variables d .

In case the disturbance variables d are absent this problem comes down to the usual system-theoretic notion of *state space equivalence*, under the *additional assumption* that the systems have *minimal state space dimension*, see e.g. [5, 6]. At the same time, the problem bears much similarity with the notion of *bisimulation* as introduced by Milner and Park [15, 21] in the study of *concurrent processes*; especially in the presence of disturbance inputs d . Finally, the notion is closely linked to the notion of *controlled invariance*, as introduced in linear systems theory by Wonham & Morse [31] and Basile & Marro [4]. The last two connections were already explored by Pappas and co-authors [19, 18, 20, 29] in the more restricted context when a system is bisimilar to an "abstraction" of itself. (Note furthermore

that in [18, 19, 29, 20] the input term Bu plays the same role as the disturbance term Gd in our setting, and thus bisimulation only involves equality of the outputs $y = Cx$.) We shall make explicit how our results generalize the results of [19, 18, 20, 29, 9] in Section 5.

The reason why the above notion of equivalence is similar to the notion of bisimulation as used in computer science can be briefly explained as follows. The notion of bisimulation has been introduced in order to handle external equivalence of (interactive) concurrent processes. The concept is especially powerful for concurrent processes which are *non-deterministic* in the sense that branching in the (discrete) state may occur while the traces (‘words’) generated by the transition system are the same. (Thus ‘non-deterministic’ is *not* meant in a stochastic sense but in the sense that the current state together with the label of the transition does not uniquely determine the next state.) In fact, the existence of a bisimulation relation between two *deterministic* processes is equivalent to equality of their external behaviors (the set of ‘traces’, or the ‘language’ generated by the process), and in this case bisimulation provides an efficient way to check equality of external behavior. For non-deterministic processes, however, bisimulation provides a finer equivalence than equality of external behavior, and, for example, also captures the *deadlock behavior* of concurrent processes. A similar picture appears to arise for bisimulation of continuous dynamical systems (1). First, a type of ‘non-determinism’ is present in systems (1) if we consider u and y as the external variables of the system (analogously to the labels of the discrete transitions of a process), while d denotes a generator for non-determinism in the evolution of the state x . If d is absent then (1) reduces to an ordinary “deterministic” system. The interpretation of d as an internal generator for non-determinism may be stressed (but for simplicity of notation we will not do so) by suppressing the appearance of the variable d by rewriting (1) equivalently as the *differential inclusion*

$$\begin{aligned} \dot{x} - Ax - Bu &\in \text{im } G \\ y &= Cx \end{aligned} \tag{2}$$

One of the results established in this paper is that for *deterministic* systems (1) (d absent) bisimulation is equivalent to equality of external behavior, while generalizing the usual notion of state space equivalence by allowing for systems with *non-minimal* state space dimension. For non-deterministic systems, however, bisimulation will be a stronger (finer) type of equivalence than equality of external behavior.

The structure of the paper is as follows. In Section 2 a (linear-)algebraic characterization of bisimulation is given, based on geometric control theory. The maximal bisimulation relation is computed in Section 3. Then Section 4 deals with reduction of dynamical systems using the notion of a bisimulation relation between the system and itself. Simulation and abstraction is treated in Section 5, making explicit the relation with previously obtained results in [19]. In Section 6 the theory is specialized to deterministic systems recovering classical results on state space equivalence and reduction. Although most of the paper is devoted to *linear* systems (1) we briefly describe in Section 7 how the theory developed for linear dynamical systems can be generalized to a *nonlinear* context by using tools from *nonlinear geometric control theory*; especially the notions of a controlled invariant submanifold and a controlled invariant distribution or foliation. Finally, Section 8 contains the conclusions and questions for further research.

2 Bisimilar linear dynamical systems

Consider two dynamical systems of the form (1):

$$\begin{aligned} \dot{x}_i &= A_i x_i + B_i u_i + G_i d_i, & x_i \in \mathcal{X}_i, u_i \in \mathcal{U}, d_i \in \mathcal{D}_i \\ \Sigma_i : & \\ y_i &= C_i x_i, & y_i \in \mathcal{Y} \quad i = 1, 2 \end{aligned} \tag{3}$$

with $\mathcal{X}_i, \mathcal{D}_i, \mathcal{U}, \mathcal{Y}$ finite-dimensional linear spaces (over \mathbb{R}). Before defining bisimulation we need to specify the solution trajectories of the systems (the ‘*semantics*’). That is, we have to specify the function classes of admissible input functions $u : [0, \infty) \rightarrow \mathcal{U}$ and admissible disturbance functions $d : [0, \infty) \rightarrow \mathcal{D}$, together with compatible function classes of state and output solutions $x : [0, \infty) \rightarrow \mathcal{X}$ and $y : [0, \infty) \rightarrow \mathcal{Y}$. For compactness of notation we will usually denote these time-functions respectively by $u(\cdot), d(\cdot), x(\cdot)$ and $y(\cdot)$. The exact class from which the functions are chosen is for purposes of this paper not really important, as long as the state trajectories $x(\cdot)$ are at least absolutely continuous. For example, we can take all functions to be C^∞ , or piecewise C^∞ with the additional requirement that $x(\cdot)$ is continuous.

Definition 2.1. A bisimulation relation between two linear systems Σ_1 and Σ_2 is a linear subspace

$$\mathcal{R} \subset \mathcal{X}_1 \times \mathcal{X}_2$$

with the following property. Take any $(x_{10}, x_{20}) \in \mathcal{R}$ and any joint input function $u_1(\cdot) = u_2(\cdot)$. Then for every disturbance function $d_1(\cdot)$ there should exist a disturbance function $d_2(\cdot)$ such that the resulting state solution trajectories $x_1(\cdot)$, with $x_1(0) = x_{10}$, and $x_2(\cdot)$, with $x_2(0) = x_{20}$, satisfy

$$(i) \quad (x_1(t), x_2(t)) \in \mathcal{R}, \quad \text{for all } t \geq 0 \tag{4}$$

$$(ii) \quad C_1 x_1(t) = C_2 x_2(t), \quad \text{for all } t \geq 0 \tag{5}$$

(or more precisely, for all $t \geq 0$ for which the trajectories are defined), and conversely, for every disturbance function $d_2(\cdot)$ there should exist a disturbance function $d_1(\cdot)$ such that again the resulting state trajectories $x_1(\cdot)$ and $x_2(\cdot)$ satisfy (4) and (5).

Hence for every pair $(x_{10}, x_{20}) \in \mathcal{R}$ all possible trajectories $x_1(\cdot)$ with $x_1(0) = x_{10}$ can be ‘simulated’ by a trajectory $x_2(\cdot)$ with $x_2(0) = x_{20}$ and conversely all possible trajectories $x_2(\cdot)$ with $x_2(0) = x_{20}$ can be ‘simulated’ by a trajectory $x_1(\cdot)$ with $x_1(0) = x_{10}$, in the sense of giving the same input-output data for all future times while $(x_1(t), x_2(t)) \in \mathcal{R}$ for all $t \geq 0$.

Remark 2.2. For linear systems it is natural to restrict to bisimulation relations \mathcal{R} which are linear subspaces. This can be done without much loss of generality. Indeed, the maximal bisimulation relation as produced by Algorithm 3.3 is a linear subspace. Furthermore, if a general subset \mathcal{R} satisfies the conditions of Definition 2.1 then so does its linear closure.

Remark 2.3. A similar definition (in the nonlinear case, cf. Section 7, and for the case that u_i is absent) was given in [9].

Remark 2.4. Note that the existence of a bisimulation relation between systems defines an *equivalence relation* between systems. Clearly $\mathcal{R}_{id} := \{(x_1, x_1) \mid x_1 \in \mathcal{X}_1\}$ is a bisimulation relation between Σ_1 given by (3) and itself. Furthermore, the existence of a bisimulation relation between Σ_1 and Σ_2 is symmetric. Finally, if $\mathcal{R}_{12} \subset \mathcal{X}_1 \times \mathcal{X}_2$ is a bisimulation relation between Σ_1 and Σ_2 , and $\mathcal{R}_{23} \subset \mathcal{X}_2 \times \mathcal{X}_3$ between Σ_2 and Σ_3 then $\mathcal{R}_{13} := \{(x_1, x_3) \mid \exists x_2 \in \mathcal{X}_2 \text{ s.t. } (x_1, x_2) \in \mathcal{R}_{12}, (x_2, x_3) \in \mathcal{R}_{23}\}$ is a bisimulation relation between Σ_1 and Σ_3 .

Definition 2.5. Two systems Σ_1 and Σ_2 as in (3) are *bisimilar*, denoted $\Sigma_1 \sim \Sigma_2$, if there exists a bisimulation relation $\mathcal{R} \subset \mathcal{X}_1 \times \mathcal{X}_2$ with the property that

$$\pi_1(\mathcal{R}) = \mathcal{X}_1, \quad \pi_2(\mathcal{R}) = \mathcal{X}_2 \quad (6)$$

where $\pi_i : \mathcal{X}_1 \times \mathcal{X}_2 \rightarrow \mathcal{X}_i$, $i = 1, 2$, denote the canonical projections.

Remark 2.6. Actually Definition 2.5 constitutes a slight departure from the definition of bisimulation relation as usually given for discrete processes [15, 21], by imposing the extra requirement (6). The reason is that in computer science discrete processes are usually defined with respect to a *fixed* initial state (or, a subset of initial conditions). In our setting we consider the behavior of the systems Σ_i for *arbitrary* initial states. Hence for every initial condition x_{10} of Σ_1 there should exist an initial condition x_{20} of Σ_2 with $(x_{10}, x_{20}) \in \mathcal{R}$ and vice versa; thus implying (6). The generalization to *subsets* of initial conditions $\mathcal{X}_{i0} \subset \mathcal{X}_i$ obviously can be done by relaxing (6) to $\pi_i(\mathcal{R}) = \mathcal{X}_{i0}$, $i = 1, 2$.

Remark 2.7. Clearly (cf. Remark 2.4), the relation \sim is an equivalence relation.

Remark 2.8. For $G_1 = G_2 = 0$ the above notion of bisimilarity is close to the usual notion of *state space equivalence* of two input-state-output systems (cf. [5, 6])

$$\begin{aligned} \Sigma_i : \quad \dot{x}_i &= A_i x_i + B_i u_i, \quad x_i \in \mathcal{X}_i, u_i \in \mathcal{U}, y_i \in \mathcal{Y} \\ y_i &= C_i x_i \quad \quad \quad i = 1, 2 \end{aligned} \quad (7)$$

Indeed, in this case one usually starts with a linear equivalence *mapping*

$$S : \mathcal{X}_1 \rightarrow \mathcal{X}_2 \quad (8)$$

which is assumed to be invertible (implying that $\dim \mathcal{X}_1 = \dim \mathcal{X}_2$) with the property that

$$Sx_1(t) = x_2(t), \quad \text{for all } t \geq 0 \quad (9)$$

$$C_1 x_1(t) = C_2 x_2(t), \quad \text{for all } t \geq 0 \quad (10)$$

for all state trajectories $x_1(\cdot)$ and $x_2(\cdot)$ resulting from initial conditions x_{10} and x_{20} related by $Sx_{10} = x_{20}$ and all input-functions $u_1(\cdot) = u_2(\cdot)$. Defining the linear subspace

$$\mathcal{R} = \{(x_1, x_2) \in \mathcal{X}_1 \times \mathcal{X}_2 \mid x_2 = Sx_1\} \quad (11)$$

(i.e., the graph of the mapping S) it is easily seen that \mathcal{R} is a bisimulation relation which satisfies $\pi_1(\mathcal{R}) = \mathcal{X}_1$ trivially and $\pi_2(\mathcal{R}) = \mathcal{X}_2$ because of invertibility of S .

Clearly, by allowing \mathcal{R} to be a *relation* instead of the graph of a mapping, the notion of bisimilarity even in the case $G_1 = G_2 = 0$ is more general than state space equivalence. In particular, we may allow \mathcal{X}_1 and \mathcal{X}_2 to be of *different dimension*. Furthermore, by doing so we incorporate in the notion of bisimilarity the notion of *reduction* of an input-state-output system to a lower-dimensional input-state-output system, and especially the reduction to a *minimal* input-state-output system, as we will see later on in Section 6.

Using ideas from the theory of controlled invariance (or (A, B) -invariance), see e.g. [31, 4], it is relatively straightforward to derive an algebraic characterization of the notion of a bisimulation relation. We do this in two steps.

Proposition 2.9. *A subspace $R \subset \mathcal{X}_1 \times \mathcal{X}_2$ is a bisimulation relation between Σ_1 and Σ_2 if and only if for all $(x_1, x_2) \in \mathcal{R}$ and all $u \in \mathcal{U}$ the following properties hold:*

(i) *For all $d_1 \in \mathcal{D}_1$ there should exist a $d_2 \in \mathcal{D}_2$ such that*

$$(A_1x_1 + B_1u + G_1d_1, A_2x_2 + B_2u + G_2d_2) \in \mathcal{R}, \quad (12)$$

and conversely for every $d_2 \in \mathcal{D}_2$ there should exist a $d_1 \in \mathcal{D}_1$ such that (12) holds.

(ii)

$$C_1x_1 = C_2x_2 \quad (13)$$

Proof. (Sketch) Consider (4). Then by differentiating $x_1(t)$ and $x_2(t)$ with respect to t and evaluating at any t we obtain (12), with $x_1 = x_1(t), x_2 = x_2(t), u = u_1(t) = u_2(t), d_1 = d_1(t), d_2 = d_2(t)$. Conversely, if (12) holds then $(\dot{x}_1(t), \dot{x}_2(t)) \in \mathcal{R}$ for all $t \geq 0$ for which the derivative exists, thus implying (4). Equivalence of (5) and (13) is obvious. ■

Theorem 2.10. *A subspace $\mathcal{R} \subset \mathcal{X}_1 \times \mathcal{X}_2$ is a bisimulation relation between Σ_1 and Σ_2 if and only if*

$$\begin{aligned} (a) \quad & \mathcal{R} + \text{im} \begin{bmatrix} G_1 \\ 0 \end{bmatrix} = \mathcal{R} + \text{im} \begin{bmatrix} 0 \\ G_2 \end{bmatrix} =: \mathcal{R}_e \\ (b) \quad & \begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix} \mathcal{R} \subset \mathcal{R}_e \\ (c) \quad & \text{im} \begin{bmatrix} B_1 \\ B_2 \end{bmatrix} \subset \mathcal{R}_e \\ (d) \quad & \mathcal{R} \subset \ker \begin{bmatrix} C_1 \\ -C_2 \end{bmatrix} \end{aligned} \quad (14)$$

Proof. Clearly (13) is equivalent with (14d). Let \mathcal{R} satisfy property (i) of Proposition 2.9. Take $u = 0$. Then for all $(x_1, x_2) \in \mathcal{R}$, and for all d_1 there exists a d_2 such that

$$(A_1x_1 + G_1d_1, A_2x_2 + G_2d_2) \in \mathcal{R} \quad (15)$$

This is a version of the so-called Modified Disturbance Decoupling Problem (with d_1 being the ‘disturbance’, which is assumed to be available for feedforward control action, and d_2 being the ‘control’, [31]). It follows from the solution of this problem (see [31]) that (15) is equivalent to

$$\begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix} \mathcal{R} \subset \mathcal{R} + \text{im} \begin{bmatrix} 0 \\ G_2 \end{bmatrix} \quad (16)$$

$$\text{im} \begin{bmatrix} G_1 \\ 0 \end{bmatrix} \subset \mathcal{R} + \text{im} \begin{bmatrix} 0 \\ G_2 \end{bmatrix}$$

Analogously, property (i) of Proposition 2.9 implies that for all $(x_1, x_2) \in \mathcal{R}$, and all d_2 there exists d_1 such that (15) holds, implying similarly

$$\begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix} \mathcal{R} \subset \mathcal{R} + \text{im} \begin{bmatrix} G_1 \\ 0 \end{bmatrix} \quad (17)$$

$$\text{im} \begin{bmatrix} 0 \\ G_2 \end{bmatrix} \subset \mathcal{R} + \text{im} \begin{bmatrix} G_1 \\ 0 \end{bmatrix}$$

The second lines of (16) and (17) are readily seen to be equivalent to (14a). Consequently, the first lines of (16) and (17) are equivalent with (14b).

Finally, consider property (i) for $x_1 = x_2 = 0$ and $d_1 = 0$. Then (12) implies

$$\text{im} \begin{bmatrix} B_1 \\ B_2 \end{bmatrix} \subset \mathcal{R} + \text{im} \begin{bmatrix} 0 \\ G_2 \end{bmatrix} = \mathcal{R}_e$$

(the same result follows by taking $d_2 = 0$ instead of $d_1 = 0$), and thus (14c). By linearity it follows that conversely (14a,b,c) imply property (i). \blacksquare

Remark 2.11. Although we have invoked in the proof of Theorem 2.10 the solution of the Modified Disturbance Decoupling Problem it is important to stress that the *interpretation* in the current case is completely different. Indeed, for bisimulation we do *not* aim to construct a feedback controller in d_2 dealing with the disturbance d_1 , and vice versa. Instead, we only derive algebraic conditions which ensure that the trajectories of Σ_1 are replicated by Σ_2 and conversely.

Remark 2.12. Note that a subspace $\mathcal{R} \subset \mathcal{X}_1 \times \mathcal{X}_2$ satisfies properties (14a,b) if and only if the mapping F (from subspaces $\mathcal{S} \subset \mathcal{X}_1 \times \mathcal{X}_2$ to subspaces $F(\mathcal{S}) \subset \mathcal{X}_1 \times \mathcal{X}_2$) defined by

$$\mathcal{S} \xrightarrow{F} \left\{ z \in \mathcal{X}_1 \times \mathcal{X}_2 \mid \begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix} z + \text{im} \begin{bmatrix} G_1 \\ 0 \end{bmatrix} \subset \mathcal{S} + \text{im} \begin{bmatrix} 0 \\ G_2 \end{bmatrix}, \right. \\ \left. \begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix} z + \text{im} \begin{bmatrix} 0 \\ G_2 \end{bmatrix} \subset \mathcal{S} + \text{im} \begin{bmatrix} G_1 \\ 0 \end{bmatrix} \right\}$$

satisfies $\mathcal{R} \subset F(\mathcal{R})$. This will be instrumental to compute the maximal bisimulation relation, see Algorithm 3.3; in fact, the maximal bisimulation relation turns out to be a *fixed-point* of this mapping. (This is well-known in the theory of bisimulation for concurrent processes [15].)

Remark 2.13. All notions and results regarding bisimulation developed in this paper for continuous-time linear dynamical systems (1) have a direct analogue for discrete-time linear systems $x(k+1) = Ax(k) + Bu(k) + Gd(k)$, $y(k) = Cx(k)$, $k \in \mathbb{Z}$. In particular, bisimulation between two discrete-time linear systems is characterized by the same conditions (14) as in Theorem 2.10.

Bisimilarity is easily seen to imply *equality of external behavior*. Consider two systems Σ_i , $i = 1, 2$, as in (3), with external behavior \mathcal{B}_i defined as

$$\mathcal{B}_i := \{(u_i(\cdot), y_i(\cdot)) \mid \exists x_i(\cdot), d_i(\cdot) \text{ such that (3) is satisfied}\} \quad (18)$$

Proposition 2.14. *Let Σ_i , $i = 1, 2$, be bisimilar. Then their external behaviors \mathcal{B}_i are equal.*

However, in the case of non-deterministic systems, that is, d_i is present, systems may have the same external behavior, while *not* being bisimilar. This is illustrated by the following simple example.

Example 2.15. Consider the two systems

$$\Sigma_1 : \begin{aligned} \dot{x}^1 &= x^2 \\ \dot{x}^2 &= d_1 \\ y_1 &= x^1 \end{aligned} \quad (19)$$

and

$$\Sigma_2 : \begin{array}{l} \dot{z} = d_2 \\ y_2 = z \end{array} \quad (20)$$

It can be readily seen that there does not exist any bisimulation relation between Σ_1 and Σ_2 (consider condition (14a)). On the other hand, if we restrict to C^∞ external behaviors then obviously $\mathcal{B}_1 = \mathcal{B}_2$. (Note the different logical quantifiers in the definition of bisimilarity and in equality of external behavior. For bisimilarity there should exist for every x^1, x^2 a z such that for every d_1 there exists a d_2 with equal external trajectories and conversely, while for equality of external behavior there should exist for every x^1, x^2, d_1 a pair z, d_2 with equal external trajectories, and conversely.)

An *interpretation* of the fact that Σ_1 and Σ_2 are not bisimilar can be given as follows. Suppose we ‘test’ the system Σ_1 at some time instant $t = t_0$ in the sense of observing one of its possible external trajectories $y_1(t), t \geq t_0$. At $t = t_0$ the system Σ_1 is in a given, but unknown, initial state $(x^1(t_0), x^2(t_0))$. Hence, all possible runs $y_1(t), t \geq t_0$, starting from this fixed initial state will have a *fixed* time-derivative $\dot{y}_1(t_0) = x^2(t_0)$ at $t = t_0$. On the other hand, for Σ_2 the possible runs $y_2(t), t \geq t_0$, can have arbitrary time-derivative at $t = t_0$. Hence, Σ_1 and Σ_2 can be considered to be externally *different*.

Remark 2.16. Note that in Example 2.15 the solution concept (‘semantics’) *does* play some role. If we choose the solutions different from C^∞ , e.g. piecewise C^∞ with $x(\cdot)$ continuous, then the outputs $y_1(\cdot)$ generated by Σ_1 will have a ‘higher degree of smoothness’ than the outputs $y_2(\cdot)$ of Σ_2 , implying that $\mathcal{B}_1 \subset \mathcal{B}_2$ while the two behaviors are not exactly equal.

For deterministic systems (d_i absent) it will be shown in Section 6 that equality of external behavior *does* imply bisimilarity.

3 Computing the maximal bisimulation relation

In this section we show how to compute the *maximal* bisimulation relation $R \subset \mathcal{X}_1 \times \mathcal{X}_2$ for two linear dynamical systems Σ_1 and Σ_2 . The way to do this is very similar to the computation of the maximal controlled invariant subspace contained in a given subspace, which is the central algorithm in linear geometric control theory [31]. Furthermore, structurally the algorithm is the same as the existing algorithms to compute the maximal bisimulation relation for two discrete processes, see e.g. [11].

First we remark that the maximal bisimulation relation exists if there exists a bisimulation relation at all (contrary to e.g. the *minimal* bisimulation relation). The argument is similar to the argument showing the existence of a maximal controlled invariant subspace, and is based on the following simple observations.

Proposition 3.1. *Let $\mathcal{R}_a \subset \mathcal{X}_1 \times \mathcal{X}_2$ and $\mathcal{R}_b \subset \mathcal{X}_1 \times \mathcal{X}_2$ be bisimulation relations. Then also $\mathcal{R}_a + \mathcal{R}_b \subset \mathcal{X}_1 \times \mathcal{X}_2$ is a bisimulation relation.*

Proof. Since $\mathcal{R}_a, \mathcal{R}_b$ are bisimulation relations they satisfy properties (14). It follows that also $\mathcal{R}_a + \mathcal{R}_b$ satisfies (14), and thus is a bisimulation relation. \blacksquare

Proposition 3.2. *Given Σ_1 and Σ_2 and suppose there exists a bisimulation relation between Σ_1 and Σ_2 . Then the maximal bisimulation relation exists.*

Proof. Suppose there exists a bisimulation relation. Let \mathcal{R}^{max} be a bisimulation relation of *maximal dimension*. Take any other bisimulation relation \mathcal{R} . Then $\mathcal{R} \subset \mathcal{R}^{max}$, since otherwise $\dim(\mathcal{R} + \mathcal{R}^{max}) > \dim \mathcal{R}^{max}$ while also $\mathcal{R} + \mathcal{R}^{max}$ is a bisimulation relation; a contradiction with the maximality of dimension of \mathcal{R}^{max} . ■

The fact that the maximal bisimulation relation exists is very fortunate because in most applications we are precisely interested in this maximal one. This is already apparent from the definition of bisimilarity including the requirement (6), but the fact also shows up in maximally *reducing* a system to a bisimilar system (Section 4).

The maximal bisimulation relation \mathcal{R}^{max} can be computed in the following way, similarly to the algorithm to compute the maximal controlled invariant subspace [31]. For notational convenience we define

$$A^\times := \begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix}, \quad G_1^\times := \begin{bmatrix} G_1 \\ 0 \end{bmatrix}, \quad G_2^\times := \begin{bmatrix} 0 \\ G_2 \end{bmatrix}, \quad C^\times := \begin{bmatrix} C_1 \\ -C_2 \end{bmatrix} \quad (21)$$

Algorithm 3.3. Given two dynamical systems Σ_1 and Σ_2 . Define the following sequence $\mathcal{R}^j, j = 0, 1, 2, \dots$, of subsets of $\mathcal{X}_1 \times \mathcal{X}_2$

$$\begin{aligned} \mathcal{R}^0 &= \mathcal{X}_1 \times \mathcal{X}_2 \\ \mathcal{R}^1 &= \{z \in \mathcal{R}^0 \mid z \in \ker C^\times\} \\ \mathcal{R}^2 &= \{z \in \mathcal{R}^1 \mid A^\times z + \text{im } G_1^\times \subset \mathcal{R}^1 + \text{im } G_2^\times, A^\times z + \text{im } G_2^\times \subset \mathcal{R}^1 + \text{im } G_1^\times\} \\ &\vdots \\ \mathcal{R}^{j+1} &= \{z \in \mathcal{R}^j \mid A^\times z + \text{im } G_1^\times \subset \mathcal{R}^j + \text{im } G_2^\times, A^\times z + \text{im } G_2^\times \subset \mathcal{R}^j + \text{im } G_1^\times\} \end{aligned} \quad (22)$$

Theorem 3.4. The sequence of subsets $\mathcal{R}^0, \mathcal{R}^1, \dots, \mathcal{R}^j, \dots$ satisfies the following properties.

1. $\mathcal{R}^j, j \geq 0$, is a linear space or empty. Furthermore $\mathcal{R}^0 \supset \mathcal{R}^1 \supset \mathcal{R}^2 \dots \supset \mathcal{R}^j \supset \mathcal{R}^{j+1} \supset \dots$
2. There exists a finite k such that $\mathcal{R}^k = \mathcal{R}^{k+1} =: \mathcal{R}^*$ and then $\mathcal{R}^j = \mathcal{R}^*$ for all $j \geq k$.
3. \mathcal{R}^* is either empty or equals the maximal subspace of $\mathcal{X}_1 \times \mathcal{X}_2$ satisfying properties (14a,b,d) of Proposition 2.10.

Proof. The proof is very similar to the proof of the corresponding properties of the algorithm for computing the maximal controlled invariant subspace [31], and hence only a sketch will be provided.

1. This is easily shown by induction.
2. Follows from Property 1 and finite-dimensionality of the subspaces involved.
3. First we show that \mathcal{R}^* obtained in Property 2, whenever non-empty, satisfies properties (14a,b,d). Since $\mathcal{R}^* = \mathcal{R}^k = \mathcal{R}^{k+1}$ we have

$$\begin{aligned} A^\times \mathcal{R}^* + \text{im } G_1^\times &\subset \mathcal{R}^* + \text{im } G_2^\times \\ A^\times \mathcal{R}^* + \text{im } G_2^\times &\subset \mathcal{R}^* + \text{im } G_1^\times \end{aligned} \quad (23)$$

which implies

$$\text{im } G_1^\times \subset \mathcal{R}^* + \text{im } G_2^\times \quad (24)$$

$$\text{im } G_2^\times \subset \mathcal{R}^* + \text{im } G_1^\times$$

and

$$A^\times \mathcal{R}^* \subset \mathcal{R}^* + \text{im } G_2^\times, A^\times \mathcal{R}^* \subset \mathcal{R}^* + \text{im } G_1^\times \quad (25)$$

Clearly (24) is equivalent to property (14a), while (25) yields property (14b). Furthermore $\mathcal{R}^* \subset \mathcal{R}^1 = \ker C^\times$, and thus \mathcal{R}^* satisfies property (14d). In order to show that \mathcal{R}^* , if non-empty, is the *maximal* subspace satisfying properties (14a,b,d), let \mathcal{R} be any subspace satisfying (14a,b,d). Clearly $\mathcal{R} \subset \mathcal{R}^1$, and by induction we prove $\mathcal{R} \subset \mathcal{R}^j$ for all j , implying that $\mathcal{R} \subset \mathcal{R}^*$. ■

If \mathcal{R}^* as obtained from Algorithm 3.3 is non-empty and satisfies property (14c), then it follows that \mathcal{R}^* is the *maximal bisimulation relation* \mathcal{R}^{max} between Σ_1 and Σ_2 , while if \mathcal{R}^* is empty or does *not* satisfy property (14c) then there does *not* exist any bisimulation relation between Σ_1 and Σ_2 . With regard to bisimilarity (Definition 2.5), we have the following immediate consequence.

Corollary 3.5. Σ_1 and Σ_2 are bisimilar if and only if \mathcal{R}^* satisfies Property (14c) and Equation (6).

In the previous section, cf. Proposition 2.14, we have seen that bisimilarity implies equality of external behavior. In fact, a computationally efficient way to check equality of external behavior of two systems $\Sigma_i, i = 1, 2$, may be to compute \mathcal{R}^* as above, and to check bisimilarity by Corollary 3.5. (Recall, however, that non-deterministic systems may have equal behavior while not being bisimilar, cf. Example 2.15.)

4 Reduction of systems by bisimulation

In this section we study the question how to *reduce* a linear dynamical system to a system with *lower state space dimension*, which is *bisimilar* to the original system, and in particular how to reduce the system to a bisimilar system with *minimal* state space dimension.

It turns out that this can be achieved by considering bisimulation relations between the system and *a copy of itself*. Furthermore, the reduction to a bisimilar system with minimal state space dimension can be performed by using the same algorithm as given in the previous section for computing the maximal bisimulation relation. Actually, this idea is well-known in the context of concurrent processes, see e.g. [11].

So let us consider a linear dynamical system as in (1):

$$\begin{aligned} \dot{x} &= Ax + Bu + Gd, & x \in \mathcal{X}, u \in \mathcal{U}, d \in \mathcal{D} \\ \Sigma : & & \\ y &= Cx, & y \in \mathcal{Y} \end{aligned} \quad (26)$$

with $\mathcal{X}, \mathcal{U}, \mathcal{Y}$ and \mathcal{D} finite-dimensional linear spaces. Now consider a bisimulation relation

between Σ and itself, that is, in view of Theorem 2.10, a subspace $\mathcal{R} \subset \mathcal{X} \times \mathcal{X}$ satisfying

$$\begin{aligned}
(a) \quad & \mathcal{R} + \text{im} \begin{bmatrix} G \\ 0 \end{bmatrix} = \mathcal{R} + \text{im} \begin{bmatrix} 0 \\ G \end{bmatrix} =: \mathcal{R}_e \\
(b) \quad & \begin{bmatrix} A & 0 \\ 0 & A \end{bmatrix} \mathcal{R} \subset \mathcal{R}_e \\
(c) \quad & \text{im} \begin{bmatrix} B \\ B \end{bmatrix} \subset \mathcal{R}_e \\
(d) \quad & \mathcal{R} \subset \ker \begin{bmatrix} C \\ -C \end{bmatrix}
\end{aligned} \tag{27}$$

First of all we note the following obvious fact:

Proposition 4.1. *The identity relation $\mathcal{R}_{id} = \{(x, x) \mid x \in \mathcal{X}\}$ is a bisimulation between Σ and itself.*

Every $\mathcal{R} \subset \mathcal{X} \times \mathcal{X}$ defines a *relation* on \mathcal{X} by saying that $x_a, x_b \in \mathcal{X}$ are related by \mathcal{R} if and only if $(x_a, x_b) \in \mathcal{R}$. For reduction we should restrict attention to $\mathcal{R} \subset \mathcal{X} \times \mathcal{X}$ such that the corresponding relation on \mathcal{X} is an *equivalence* relation, i.e., \mathcal{R} is reflexive ($(x, x) \in \mathcal{R}$ for all $x \in \mathcal{X}$), symmetric ($(x_a, x_b) \in \mathcal{R} \iff (x_b, x_a) \in \mathcal{R}$), and transitive ($(x_a, x_b) \in \mathcal{R}, (x_b, x_c) \in \mathcal{R} \Rightarrow (x_a, x_c) \in \mathcal{R}$). This can be done without loss of generality. Indeed, by Proposition 4.1 we may always add to any bisimulation relation \mathcal{R} the identity bisimulation relation \mathcal{R}_{id} , thus enforcing reflexivity. Furthermore, let \mathcal{R} satisfy (27), then also the inverse relation $\mathcal{R}^{-1} := \{(x_a, x_b) \mid (x_b, x_a) \in \mathcal{R}\}$ satisfies (27), implying that the symmetric closure $\mathcal{R} + \mathcal{R}^{-1}$ satisfies (27). Finally, for *linear* relations reflexivity and symmetry already *implies* transitivity: if $(x_a, x_b), (x_b, x_c) \in \mathcal{R}$, then $(x_a - x_c, 0) = (x_a, x_b) - (x_c, x_b) \in \mathcal{R}$, and thus $(x_a, x_c) = (x_a - x_c, 0) + (x_c, x_c) \in \mathcal{R}$. Any equivalence relation $\mathcal{R} \subset \mathcal{X} \times \mathcal{X}$ can be uniquely associated with a linear subspace $\bar{\mathcal{R}} \subset \mathcal{X}$ defined as follows:

$$\bar{\mathcal{R}} := \{x_a - x_b \mid (x_a, x_b) \in \mathcal{R}\} \tag{28}$$

Indeed, $\bar{\mathcal{R}}$ defined by (28) is a *linear space* if and only if \mathcal{R} is reflexive and symmetric (and therefore an equivalence relation). Reduction can now be performed by factoring out \mathcal{X} by the subspace $\bar{\mathcal{R}}$, thereby obtaining the linear space of equivalence classes of the equivalence relation \mathcal{R} . In terms of the subspace $\bar{\mathcal{R}}$ conditions (27) reduce as follows.

Theorem 4.2. *Let $\mathcal{R} \subset \mathcal{X} \times \mathcal{X}$ be an equivalence relation, and define $\bar{\mathcal{R}} \subset \mathcal{X}$ as in (28). Conditions (27a,b,c,d) for \mathcal{R} are equivalent to*

$$\begin{aligned}
A\bar{\mathcal{R}} & \subset \bar{\mathcal{R}} + \text{im } G \\
\bar{\mathcal{R}} & \subset \ker C
\end{aligned} \tag{29}$$

Proof. It is readily seen that (27b,d) are equivalent to (29). Satisfaction of (27a,c) follows from reflexivity of \mathcal{R} . ■

A subspace $\bar{\mathcal{R}}$ satisfying the first line of (29) is called a *controlled invariant subspace* (with respect to A and $\text{im } G$), cf.[4, 31]. Thus there is a one-to-one correspondence between bisimulation equivalence relations \mathcal{R} and controlled invariant subspaces $\bar{\mathcal{R}}$ contained in $\ker C$.

As a consequence of Proposition 4.1 the *maximal* bisimulation $\mathcal{R}^{max} = \mathcal{R}^*$ between Σ and itself exists, and contains \mathcal{R}_{id} . Hence \mathcal{R}^* is reflexive, while by symmetry of the data it follows that the symmetric closure of \mathcal{R}^* (adjoining (x_b, x_a) if $(x_a, x_b) \in \mathcal{R}^*$) also satisfies (27a,b,d), and hence \mathcal{R}^* is symmetric. Thus the maximal bisimulation relation \mathcal{R}^* is an equivalence relation. The corresponding subspace $\bar{\mathcal{R}}^* \subset \mathcal{X}$ is the *maximal* controlled invariant subspace contained in $\ker C$.

It is now clear how to reduce Σ to a lower-dimensional system that is bisimilar to Σ . Let R be a bisimulation equivalence relation. Define the reduced state space

$$\mathcal{X}_{\mathcal{R}} := \mathcal{X}/\bar{\mathcal{R}} \quad (30)$$

with canonical projection $\Pi_{\mathcal{R}} : \mathcal{X} \rightarrow \mathcal{X}/\bar{\mathcal{R}}$. By the first line of (29) there exists a "feedback" map K such that

$$(A + GK)\bar{\mathcal{R}} \subset \bar{\mathcal{R}} \quad (31)$$

and thus $A + GK$ projects to a linear map

$$A_{\mathcal{R}} : \mathcal{X}_{\mathcal{R}} \rightarrow \mathcal{X}_{\mathcal{R}} \quad (32)$$

satisfying $A_{\mathcal{R}}\Pi_{\mathcal{R}} = \Pi_{\mathcal{R}}(A + GK)$. Furthermore, define

$$G_{\mathcal{R}} := \Pi_{\mathcal{R}}G, \quad B_{\mathcal{R}} := \Pi_{\mathcal{R}}B, \quad (33)$$

and by the second line of (29) we may define

$$C_{\mathcal{R}} : \mathcal{X}_{\mathcal{R}} \rightarrow \mathcal{Y} \quad (34)$$

such that $C_{\mathcal{R}}\Pi_{\mathcal{R}} = C$. Together this defines a reduced system

$$\begin{aligned} \dot{x}_{\mathcal{R}} &= A_{\mathcal{R}}x_{\mathcal{R}} + B_{\mathcal{R}}u + G_{\mathcal{R}}d \\ \Sigma_{\mathcal{R}} : \quad y &= C_{\mathcal{R}}x_{\mathcal{R}} \end{aligned} \quad (35)$$

Proposition 4.3. *Let \mathcal{R} be a bisimulation equivalence relation between Σ and itself, and construct $\Sigma_{\mathcal{R}}$ as above. Then $\Sigma_{\mathcal{R}}$ is bisimilar to Σ . Furthermore, let \mathcal{R}^* denote the maximal bisimulation relation between Σ and itself. Then $\Sigma_{\mathcal{R}^*}$ is the smallest system that is bisimilar by reduction to Σ .*

Proof. Define the following relation $\mathcal{S}_{\mathcal{R}} \subset \mathcal{X} \times \mathcal{X}_{\mathcal{R}}$

$$(x, x_{\mathcal{R}}) \in \mathcal{S}_{\mathcal{R}} \iff x_{\mathcal{R}} = \Pi_{\mathcal{R}}(x) \quad (36)$$

This is readily seen to be a bisimulation relation which obviously satisfies (6). ■

The idea of *reducing* a linear dynamical system by factorizing the state space \mathcal{X} by the subspace corresponding to a bisimulation equivalence relation can be put into a more general

context. Indeed, consider a bisimulation relation $\mathcal{R} \subset \mathcal{X}_1 \times \mathcal{X}_2$ between two systems $\Sigma_i, i = 1, 2$, as given in (3). Then define the following subspace of \mathcal{X}_1 , respectively \mathcal{X}_2 :

$$\begin{aligned}\bar{\mathcal{R}}_1 &:= \{x_1 \in \mathcal{X}_1 \mid (x_1, 0) \in \mathcal{R}\} \\ \bar{\mathcal{R}}_2 &:= \{x_2 \in \mathcal{X}_2 \mid (0, x_2) \in \mathcal{R}\}\end{aligned}\tag{37}$$

Since \mathcal{R} is a bisimulation relation it immediately follows from (14a,b,d) that

$$\begin{aligned}A_1\bar{\mathcal{R}}_1 &\subset \bar{\mathcal{R}}_1 + \text{im } G_1, & \bar{\mathcal{R}}_1 &\subset \ker C_1 \\ A_2\bar{\mathcal{R}}_2 &\subset \bar{\mathcal{R}}_2 + \text{im } G_2, & \bar{\mathcal{R}}_2 &\subset \ker C_2\end{aligned}\tag{38}$$

Hence, similarly as above, the systems Σ_i for $i = 1, 2$ can be reduced (via bisimulation) to systems

$$\begin{aligned}\Sigma_{\mathcal{R}_i} : \quad \dot{x}_{\mathcal{R}_i} &= A_{\mathcal{R}_i}x_{\mathcal{R}_i} + B_{\mathcal{R}_i}u + G_{\mathcal{R}_i}d \\ y &= C_{\mathcal{R}_i}x_{\mathcal{R}_i}\end{aligned}\tag{39}$$

living on the state spaces $\mathcal{X}_i/\bar{\mathcal{R}}_i, i = 1, 2$, with canonical projections $\Pi_{\mathcal{R}_i} : \mathcal{X}_i \rightarrow \mathcal{X}_i/\bar{\mathcal{R}}_i, i = 1, 2$. Furthermore, the relation $\mathcal{R} \subset \mathcal{X}_1 \times \mathcal{X}_2$ reduces to a relation $\tilde{\mathcal{R}} \subset \mathcal{X}_1/\bar{\mathcal{R}}_1 \times \mathcal{X}_2/\bar{\mathcal{R}}_2$ defined by

$$\tilde{\mathcal{R}} := \{(x_{\mathcal{R}_1}, x_{\mathcal{R}_2}) \mid (x_1, x_2) \in \mathcal{R} \text{ for all } (x_1, x_2) \text{ with } \Pi_{\mathcal{R}_1}x_1 = x_{\mathcal{R}_1}, \Pi_{\mathcal{R}_1}x_2 = x_{\mathcal{R}_2}\}\tag{40}$$

Moreover, it is straightforward to check that $\tilde{\mathcal{R}}$ defines a *bisimulation* between $\Sigma_{\mathcal{R}_1}$ and $\Sigma_{\mathcal{R}_2}$. In fact we have the following

Theorem 4.4. *Let $\mathcal{R} \subset \mathcal{X}_1 \times \mathcal{X}_2$ be bisimulation relation between two systems $\Sigma_i, i = 1, 2$. Then the systems Σ_i can be reduced by bisimulation to the systems $\Sigma_{\mathcal{R}_i}, i = 1, 2$, with $\tilde{\mathcal{R}}$ as defined in (40) a bisimulation relation between them. This bisimulation relation $\tilde{\mathcal{R}}$ is one-to-one in the sense that $(x_{\mathcal{R}_1}, x_{\mathcal{R}_2}), (x'_{\mathcal{R}_1}, x_{\mathcal{R}_2}), (x_{\mathcal{R}_1}, x'_{\mathcal{R}_2}) \in \tilde{\mathcal{R}}$ implies $x_{\mathcal{R}_1} = x'_{\mathcal{R}_1}$ and $x_{\mathcal{R}_2} = x'_{\mathcal{R}_2}$. Furthermore, $\Sigma_i, i = 1, 2$, are bisimilar with bisimulation relation \mathcal{R} if and only if $\Sigma_{\mathcal{R}_i}, i = 1, 2$ are bisimilar with bisimulation relation $\tilde{\mathcal{R}}$. In this case the bisimulation relation $\tilde{\mathcal{R}}$ corresponds to an invertible map $S : \mathcal{X}_1/\bar{\mathcal{R}}_1 \rightarrow \mathcal{X}_2/\bar{\mathcal{R}}_2$.*

Theorem 4.4 expresses that every bisimulation relation can be factorized into a one-to-one bisimulation relation between reduced systems, which are themselves bisimilar (by reduction) to the original systems. Furthermore, if $\Sigma_i, i = 1, 2$, are bisimilar then their reduced systems are linked by an *invertible mapping* (a state space equivalence mapping).

Corollary 4.5. *Suppose the maximal bisimulation relation \mathcal{R}^{max} between Σ_1 and Σ_2 exists and satisfies $\bar{\mathcal{R}}_1^{max} = 0, \bar{\mathcal{R}}_2^{max} = 0$ and $\pi_1(\mathcal{R}^{max}) = \mathcal{X}_1, \pi_2(\mathcal{R}^{max}) = \mathcal{X}_2$. Then $\mathcal{R}^{max} = \{(x_1, x_2 = Sx_1) \mid x_1 \in \mathcal{X}_1\}$ for some invertible map $S : \mathcal{X}_1 \rightarrow \mathcal{X}_2$.*

5 Simulation and abstraction

A one-sided version of the notions of bisimulation relation and bisimilarity as provided in Definitions 2.1 and 2.5 can be stated as follows:

Definition 5.1. Consider Σ_1 and Σ_2 as given in (3). A *simulation relation* of Σ_1 by Σ_2 is a linear subspace

$$\mathcal{S} \subset \mathcal{X}_1 \times \mathcal{X}_2$$

with the following property. Take any $(x_{10}, x_{20}) \in \mathcal{S}$ and any joint input function $u_1(\cdot) = u_2(\cdot)$. Then for every disturbance function $d_1(\cdot)$ there should exist a disturbance function $d_2(\cdot)$ such that the resulting state solution trajectories $x_1(\cdot)$, with $x_1(0) = x_{10}$, and $x_2(\cdot)$, with $x_2(0) = x_{20}$, satisfy

$$(i) \quad (x_1(t), x_2(t)) \in \mathcal{S}, \quad \text{for all } t \geq 0 \quad (41)$$

$$(ii) \quad C_1 x_1(t) = C_2 x_2(t), \quad \text{for all } t \geq 0 \quad (42)$$

Furthermore, Σ_1 is *simulated* by Σ_2 if the simulation relation \mathcal{S} satisfies $\pi_1(\mathcal{S}) = \mathcal{X}_1$ with π_1 the canonical projection of $\mathcal{X}_1 \times \mathcal{X}_2$ to \mathcal{X}_1 .

For several purposes the existence of a simulation relation instead of a bisimulation relation is already sufficient. One may for example think of a system description capturing the required specifications on a system to be designed: if the actual closed-loop system is simulated by the specified system then at least one knows that there is no unwanted behavior in the actual closed-loop system. Or, one may ‘approximate’ a given system by a lower-dimensional system in the sense that the given system is simulated by the approximating system. (This is the idea of *abstraction* as will be discussed later on.) Then checking certain properties for the approximating system will also guarantee these properties for the given system.

By a ‘one-sided’ version of Theorem 2.10, see especially Equation (16), we obtain

Proposition 5.2. $\mathcal{S} \subset \mathcal{X}_1 \times \mathcal{X}_2$ is a simulation relation of Σ_1 by Σ_2 if and only if

$$\begin{aligned} (a) \quad & \mathcal{S} + \text{im} \begin{bmatrix} G_1 \\ 0 \end{bmatrix} \subset \mathcal{S} + \text{im} \begin{bmatrix} 0 \\ G_2 \end{bmatrix} \\ (b) \quad & \begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix} \mathcal{S} \subset \mathcal{S} + \text{im} \begin{bmatrix} 0 \\ G_2 \end{bmatrix} \\ (c) \quad & \text{im} \begin{bmatrix} B_1 \\ B_2 \end{bmatrix} \subset \mathcal{S} + \text{im} \begin{bmatrix} 0 \\ G_2 \end{bmatrix} \\ (d) \quad & \mathcal{S} \subset \ker \begin{bmatrix} C_1 \\ -C_2 \end{bmatrix} \end{aligned} \quad (43)$$

For discrete processes it may occur that there is a simulation relation of Σ_1 by Σ_2 and a simulation relation of Σ_2 by Σ_1 while there does *not* exist a bisimulation relation between Σ_1 and Σ_2 , cf. [16, 7]. In the present context, however, we have the following fact. (Recall the definition of the inverse relation $\mathcal{T}^{-1} := \{(x_a, x_b) \mid (x_b, x_a) \in \mathcal{T}\}$.)

Proposition 5.3. Let $\mathcal{S} \subset \mathcal{X}_1 \times \mathcal{X}_2$ be a simulation relation of Σ_1 by Σ_2 and let $\mathcal{T} \subset \mathcal{X}_2 \times \mathcal{X}_1$ be a simulation relation of Σ_2 by Σ_1 . Then $\mathcal{R} := \mathcal{S} + \mathcal{T}^{-1}$ is a bisimulation relation between Σ_1 and Σ_2 .

Proof. Let \mathcal{S} satisfy (43) and let \mathcal{T} satisfy (43) with index 1 throughout replaced by 2. It follows that

$$\mathcal{T}^{-1} + \mathcal{S} + \text{im} \begin{bmatrix} G_1 \\ 0 \end{bmatrix} \subset \mathcal{S} + \mathcal{T}^{-1} + \text{im} \begin{bmatrix} 0 \\ G_2 \end{bmatrix} \subset \mathcal{S} + \mathcal{T}^{-1} + \text{im} \begin{bmatrix} G_1 \\ 0 \end{bmatrix}$$

and thus (14a) results for $\mathcal{R} = \mathcal{S} + \mathcal{T}^{-1}$. Similarly, \mathcal{R} satisfies (14b,c,d). ■

An interesting consequence of Proposition 5.3 is the following. First we note that if there exists a simulation relation \mathcal{S} of Σ_1 by Σ_2 then there also exists the *maximal* simulation relation \mathcal{S}^{max} of Σ_1 by Σ_2 . The argument is the same as in the case of bisimulation (cf. Section 3): if \mathcal{S}_1 and \mathcal{S}_2 are simulation relations then also $\mathcal{S}_1 + \mathcal{S}_2$ is one. Furthermore, the maximal simulation relation \mathcal{S}^{max} can be computed by the following simplified version of Algorithm 3.3. Define the decreasing sequence $\mathcal{S}^j, j = 0, 1, 2, \dots$ of linear subspaces of $\mathcal{X}_1 \times \mathcal{X}_2$ as

$$\begin{aligned}
\mathcal{S}^0 &= \mathcal{X}_1 \times \mathcal{X}_2 \\
\mathcal{S}^1 &= \{z \in \mathcal{S}^0 \mid z \in \ker C^\times\} \\
\mathcal{S}^2 &= \{z \in \mathcal{S}^1 \mid A^\times z + \text{im } G_1^\times \subset \mathcal{S}^1 + \text{im } G_2^\times\} \\
&\vdots \\
\mathcal{S}^{j+1} &= \{z \in \mathcal{S}^j \mid A^\times z + \text{im } G_1^\times \subset \mathcal{S}^j + \text{im } G_2^\times\}
\end{aligned} \tag{44}$$

Proposition 5.4. *Suppose there exists a simulation of Σ_1 by Σ_2 , and a simulation of Σ_2 by Σ_1 (or equivalently, cf. Proposition 5.3, a bisimulation relation between Σ_1 and Σ_2). Let $\mathcal{S}^{max} \subset \mathcal{X}_1 \times \mathcal{X}_2$ denote the maximal simulation relation of Σ_1 by Σ_2 , and $\mathcal{T}^{max} \subset \mathcal{X}_2 \times \mathcal{X}_1$ the maximal simulation relation of Σ_2 by Σ_1 . Then $\mathcal{S}^{max} = (\mathcal{T}^{max})^{-1} = \mathcal{R}^{max}$, with \mathcal{R}^{max} the maximal bisimulation relation.*

Proof. By Proposition 5.3 $\mathcal{S}^{max} + (\mathcal{T}^{max})^{-1}$ is a bisimulation relation, and thus also a simulation relation of Σ_1 by Σ_2 . By maximality of \mathcal{S}^{max} it follows that $\mathcal{S}^{max} + (\mathcal{T}^{max})^{-1} = \mathcal{S}^{max}$, and thus $(\mathcal{T}^{max})^{-1} \subset \mathcal{S}^{max}$. The converse inclusion holds by considering \mathcal{T}^{max} . Hence $\mathcal{S}^{max} = (\mathcal{T}^{max})^{-1}$, which is by Proposition 5.3 a bisimulation relation, and thus contained in \mathcal{R}^{max} . Since \mathcal{R}^{max} is also a simulation relation equality follows. \blacksquare

Hence, if there exists a bisimulation relation between Σ_1 and Σ_2 , then \mathcal{R}^{max} can be also computed using the simplified algorithm (44).

Now let us recall the definition of an *abstraction* as introduced in a general context in [2, 10, 18, 20], and more specifically in [19, 29]. Consider a linear system as before, cf. (26)

$$\begin{aligned}
\dot{x} &= Ax + Bu + Gd, \quad x \in \mathcal{X}, u \in \mathcal{U}, d \in \mathcal{D} \\
\Sigma : \\
y &= Cx, \quad y \in \mathcal{Y}
\end{aligned} \tag{45}$$

together with a surjective linear map $H : \mathcal{X} \rightarrow \mathcal{Z}$, \mathcal{Z} being another linear space, satisfying

$$\ker H \subset \ker C \tag{46}$$

Clearly this implies that there exists a unique linear map $\bar{C} : \mathcal{Z} \rightarrow \mathcal{Y}$ such that

$$C = \bar{C}H \tag{47}$$

Then define the following dynamical system on \mathcal{Z}

$$\begin{aligned}
\dot{z} &= \bar{A}z + \bar{B}u + \bar{G}d, \quad z \in \mathcal{Z}, u \in \mathcal{U}, d \in \mathcal{D} \\
\bar{\Sigma} : \\
y &= \bar{C}z, \quad y \in \mathcal{Y}
\end{aligned} \tag{48}$$

with

$$\begin{aligned}\bar{A} &:= HAH^+ \\ \bar{B} &:= HB \\ \bar{G} &:= [HG:HA v_1:\dots:HA v_k]\end{aligned}\tag{49}$$

where H^+ denotes the Moore-Penrose pseudo-inverse of H , and v_1, \dots, v_k span $\ker H$.

We think of $\bar{\Sigma}$ as an ‘*abstraction*’ of Σ in the sense that we factor out the part of the state variables $x \in \mathcal{X}$ corresponding to $\ker H$. This idea was put forward in a series of innovative papers by Pappas and co-authors (see especially [19]) in the slightly more restricted case where $B = 0$. (N.B., actually in [19] the ‘generator for non-determinism’ d is denoted by u .)

It has been shown in [19] that Σ is simulated by $\bar{\Sigma}$, as also follows from Proposition 5.2 by taking the simulation relation $\mathcal{R} := \{(x, z) \mid z = Hx\}$. We may think of the transformation of Σ into $\bar{\Sigma}$ as ‘lumping’ the states x corresponding to the same vector $z = Hx$. The underlying idea again comes from the theory of interactive processes, where it is often found convenient to replace a process by a process with a smaller number of states, which however replicates the trajectories of the original system (but in general contains more). Thus $\bar{\Sigma}$ can be thought of as a ‘conservative estimate’ of Σ .

A main result of [19] is that $\mathcal{R} = \{(x, z) \mid z = Hx\}$ is actually a *bisimulation* relation between Σ and $\bar{\Sigma}$ if and only if $\ker H$ is *controlled invariant* with respect to A and $\text{im } G$. This also follows directly from Theorem 4.2. Generalization of this result to the nonlinear case (see also Section 7) was obtained in [29].

6 Deterministic systems

In this section we specialize the results of the previous section to systems without disturbances d (that is, ‘deterministic systems’), and relate the results to well-known results on state space equivalence and state space reduction of standard input-state-output systems [5, 6, 22].

Thus consider two systems

$$\begin{aligned}\Sigma_i : \quad \dot{x}_i &= A_i x_i + B_i u_i, \quad x_i \in \mathcal{X}_i, u_i \in \mathcal{U} \\ y_i &= C_i x_i, \quad y_i \in \mathcal{Y}, \quad i = 1, 2\end{aligned}\tag{50}$$

with $\mathcal{X}_i, \mathcal{U}, \mathcal{Y}$ finite-dimensional linear spaces. Theorem 2.10 is readily seen to specialize to

Corollary 6.1. *A subspace $\mathcal{R} \subset \mathcal{X}_1 \times \mathcal{X}_2$ is a bisimulation relation between Σ_1 and Σ_2 given in (50) if and only if*

$$\begin{aligned}\begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix} \mathcal{R} &\subset \mathcal{R} \\ \text{im} \begin{bmatrix} B_1 \\ B_2 \end{bmatrix} &\subset \mathcal{R} \subset \ker \begin{bmatrix} C_1 \\ -C_2 \end{bmatrix}\end{aligned}\tag{51}$$

For deterministic systems Σ_i it turns out that equality of external behaviors \mathcal{B}_i , cf. (18), implies bisimilarity (contrary to the case of non-deterministic systems, see Example 2.15). This will be shown using the maximal bisimulation relation. First note that for deterministic systems \mathcal{R}^* as produced by Algorithm 3.3 is non-empty (since $\text{im } G_i^\times$ are zero subspaces).

Theorem 6.2. Consider two systems Σ_1 and Σ_2 as given in (50) with equal external behaviors $\mathcal{B}_1 = \mathcal{B}_2$. Then Σ_1 and Σ_2 are bisimilar with bisimulation relation $\mathcal{R}^{max} = \mathcal{R}^*$ as produced by Algorithm 3.3.

Proof. It is easily seen that \mathcal{R}^* as produced by Algorithm 3.3 is given as

$$\mathcal{R}^* = \ker \begin{bmatrix} C_1 & \vdots & -C_2 \\ C_1 A_1 & \vdots & -C_2 A_2 \\ & \vdots & \\ C_1 A_1^k & \vdots & -C_2 A_2^k \\ & \vdots & \end{bmatrix} \quad (52)$$

By considering initial conditions for Σ_1 and Σ_2 both equal to zero it is clear that equality of external behaviors implies

$$\text{im} \begin{bmatrix} B_1 \\ B_2 \end{bmatrix} \subset \mathcal{R}^*,$$

and thus \mathcal{R}^* satisfies (51) and defines a bisimulation relation. What remains to be checked is that $\pi_i(\mathcal{R}^*) = \mathcal{X}_i, i = 1, 2$. This follows by considering $u = 0$ and then noting that equality of external behavior of Σ_1 and Σ_2 implies that for every $x_1 \in \mathcal{X}_1$ there exists an $x_2 \in \mathcal{X}_2$ such that $C_1 e^{A_1 t} x_1 = C_2 e^{A_2 t} x_2$, and conversely. Repeated differentiation of this equality with respect to t and evaluation at $t = 0$ yields in view of (52) that $(x_1, x_2) \in \mathcal{R}^*$. ■

While equality of external behaviors of two deterministic systems has been shown to be equivalent to their bisimilarity, the *existence* of a bisimulation relation can be characterized in the following way, relating to classical concepts of system equivalence [5, 6].

Theorem 6.3. There exists a bisimulation relation \mathcal{R} between Σ_1 and Σ_2 given in (50) if and only the Markov parameters of Σ_1 and Σ_2 are equal, that is

$$C_1 A_1^k B_1 = C_2 A_2^k B_2, \quad k = 0, 1, 2, \dots, \quad (53)$$

or equivalently if their transfer matrices $G_i(s) := C_i (Is - A_i)^{-1} B_i, i = 1, 2$, are the same.

Moreover, any bisimulation relation \mathcal{R} between Σ_1 and Σ_2 satisfies

$$\pi_i(\mathcal{R}) \supset \text{Reach}(\Sigma_i) \quad (54)$$

with $\text{Reach}(\Sigma_i) = \text{im} \begin{bmatrix} B_i \vdots A_i B_i \vdots \dots \vdots A_i^k B_i \vdots \dots \end{bmatrix}$ the reachable subspace of Σ_i . Hence, if $\Sigma_i, i = 1, 2$, are controllable, then Σ_1 and Σ_2 are bisimilar if and only if $G_1(s) = G_2(s)$.

Proof. Condition (53) is equivalent to the property that the subspace

$$\text{im} \begin{bmatrix} B_1 \vdots A_1 B_1 \vdots \dots \vdots A_1^k B_1 \vdots \dots \\ B_2 \vdots A_2 B_2 \vdots \dots \vdots A_2^k B_2 \vdots \dots \end{bmatrix},$$

which is the smallest subspace of $\mathcal{X}_1 \times \mathcal{X}_2$ that is invariant under $\begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix}$ and containing $\text{im} \begin{bmatrix} B_1 \\ B_2 \end{bmatrix}$, is contained in $\ker \begin{bmatrix} C_1 \vdots -C_2 \end{bmatrix}$. Thus (53) is equivalent to the existence of $\mathcal{R} \subset \mathcal{X}_1 \times \mathcal{X}_2$ satisfying (51), while (54) holds. Controllability is equivalent to $\text{Reach}(\Sigma_i) = \mathcal{X}_i$, implying by (54) that $\pi_i(\mathcal{R}) = \mathcal{X}_i$. ■

With regard to *reduction* by bisimulation we have the following corollaries of Proposition 4.3 and Theorem 4.4:

Corollary 6.4. *Consider*

$$\begin{aligned} \dot{x} &= Ax + Bu, \quad x \in \mathcal{X}, u \in \mathcal{U} \\ \Sigma : \\ y &= Cx, \quad y \in \mathcal{Y} \end{aligned} \tag{55}$$

Then $\mathcal{R} \subset \mathcal{X} \times \mathcal{X}$ is a bisimulation equivalence relation between Σ and itself if and only if the subspace, cf. (28), $\bar{\mathcal{R}} = \{x_a - x_b \mid (x_a, x_b) \in \mathcal{R}\}$ is an unobservable subspace of the system (i.e., $A\bar{\mathcal{R}} \subset \bar{\mathcal{R}}, \bar{\mathcal{R}} \subset \ker C$). The state space \mathcal{X} of Σ can be factored out by $\bar{\mathcal{R}}$ so as to obtain $\Sigma_{\mathcal{R}}$. Furthermore $\bar{\mathcal{R}}^*$ corresponding to the maximal bisimulation relation $\mathcal{R}^{\max} = \mathcal{R}^*$ between Σ and itself equals the maximal unobservability subspace of (55), and can be factored out so as to obtain an observable system $\Sigma_{\mathcal{R}^*}$. Conversely, if Σ is observable, then $\mathcal{R}^* = \mathcal{R}_{id}$.

Corollary 6.5. *Let $\mathcal{R} \subset \mathcal{X}_1 \times \mathcal{X}_2$ be a bisimulation relation between two systems $\Sigma_i, i = 1, 2$ given by (50). Then the systems Σ_i can be reduced by bisimulation to the systems $\Sigma_{\mathcal{R}_i}, i = 1, 2$, with $\tilde{\mathcal{R}}$ as defined in (40) a bisimulation relation between them. This bisimulation relation $\tilde{\mathcal{R}}$ is one-to-one. Furthermore, $\Sigma_i, i = 1, 2$, are bisimilar with bisimulation relation \mathcal{R} if and only if $\Sigma_{\mathcal{R}_i}, i = 1, 2$, are bisimilar with bisimulation relation $\tilde{\mathcal{R}}$. In this case the bisimulation relation $\tilde{\mathcal{R}}$ corresponds to an invertible map. If Σ_1 and Σ_2 are bisimilar and both are observable, then the maximal bisimulation relation \mathcal{R}^* is the graph of an invertible map $S : \mathcal{X}_1 \rightarrow \mathcal{X}_2$ satisfying*

$$SA_1S^{-1} = A_2, SB_1 = B_2, C_1 = C_2S \tag{56}$$

Furthermore, by maximality of \mathcal{R}^* the map S satisfying (56) is unique.

Remark 6.6. From the proof of Theorem 6.2 it follows that if Σ_1 and Σ_2 are bisimilar then $(x_1, x_2) \in \mathcal{R}^*$ implies that $C_1e^{A_1t}x_1 = C_2e^{A_2t}x_2$ for all $t \geq 0$. Hence in this case \mathcal{R}^* is actually given by

$$\mathcal{R}^* = \ker \begin{bmatrix} C_1 & \vdots & -C_2 \\ C_1A_1 & \vdots & -C_2A_2 \\ \vdots & & \\ C_1A_1^{n-1} & \vdots & -C_2A_2^{n-1} \end{bmatrix} \tag{57}$$

where $n = \max(n_1, n_2), n_i = \dim \mathcal{X}_i$. From here it is directly seen that $\bar{\mathcal{R}}_i^*$ is the maximal unobservable space of Σ_i .

If Σ_1 and Σ_2 are bisimilar and at least one of the systems, say Σ_2 , is observable, then \mathcal{R}^* is the graph of the mapping $W_2^{-1}W_1 : \mathcal{X}_1 \rightarrow \mathcal{X}_2$ with

$$W_1 = \begin{bmatrix} C_1 \\ C_1A_1 \\ \vdots \\ C_1A_1^{n-1} \end{bmatrix}, W_2 = \begin{bmatrix} C_2 \\ C_2A_2 \\ \vdots \\ C_2A_2^{n-1} \end{bmatrix},$$

where $n = n_2$ and W_2^{-1} denotes the set-theoretic inverse. (See [22] for a related result in the case of scalar output systems which both are observable.)

Finally, by combining Theorem 6.3 with Corollary 6.5 we recover the classical state-space equivalence theorem for controllable and observable deterministic systems, see e.g. [5].

Corollary 6.7. *Let the systems $\Sigma_i, i = 1, 2$, given by (50) be observable and controllable. Then $\Sigma_i, i = 1, 2$, are bisimilar if and only if $G_1(s) = G_2(s)$, while the bisimulation relation is given as $\{(x_1, Sx_2) \mid x_1 \in \mathcal{X}_1\}$ for a uniquely defined invertible map $S : \mathcal{X}_1 \rightarrow \mathcal{X}_2$ (the state space equivalence map) satisfying (56).*

7 Bisimulation of nonlinear systems

In this section we briefly indicate the extension of the theory of bisimulation as established for linear dynamical systems (3) to *nonlinear* systems

$$\begin{aligned} \Sigma_i : \quad \dot{x}_i &= a_i(x_i) + b_i(x_i)u_i + g_i(x_i)d_i, \quad x_i \in \mathcal{X}, u_i \in \mathcal{U}, d_i \in \mathcal{D}_i \\ y_i &= c_i(x_i), \quad y_i \in \mathcal{Y}, \quad i = 1, 2 \end{aligned} \quad (58)$$

Here, as before, $\mathcal{D}_i, \mathcal{U}$ and \mathcal{Y} are finite-dimensional linear spaces, but the state space \mathcal{X}_i is a manifold. Furthermore, $a_i(x_i)$ and the columns of the matrices $b_i(x_i)$ and $g_i(x_i)$ denote vector fields on \mathcal{X}_i . All data are assumed to be *smooth* (C^∞). While in the linear case we restricted attention to *linear* bisimulation relations in this case it is natural to consider nonlinear bisimulation relations \mathcal{R} in Definition 2.1 which are *submanifolds* of $\mathcal{X}_1 \times \mathcal{X}_2$. (As usual, ‘submanifold’ throughout stands for ‘imbedded smooth submanifold’, see e.g. [1, 17].)

Furthermore, like in Definition 2.5, we say that Σ_1 and Σ_2 as in (58) are bisimilar if there exists a nonlinear bisimulation $\mathcal{R} \subset \mathcal{X}_1 \times \mathcal{X}_2$ with the property that $\pi_i(\mathcal{R}) = \mathcal{X}_i, i = 1, 2$.

Using the theory of controlled invariance for nonlinear systems, see e.g. [12, 17], it is immediate to derive the following generalization of Proposition 2.9.

Proposition 7.1. *A submanifold $\mathcal{R} \subset \mathcal{X}_1 \times \mathcal{X}_2$ is a bisimulation relation between Σ_1 and Σ_2 if and only if for all $(x_1, x_2) \in \mathcal{R}$ and all $u \in \mathcal{U}$ the following properties hold:*

(i) *For all $d_1 \in \mathcal{D}_1$ there should exist a $d_2 \in \mathcal{D}_2$ such that*

$$(a_1(x_1) + b_1(x_1)u + g_1(x_1)d_1, a_2(x_2) + b_2(x_2)u + g_2(x_2)d_2) \in T_{(x_1, x_2)}\mathcal{R} \quad (59)$$

and conversely for every $d_2 \in \mathcal{D}_2$ there should exist a $d_1 \in \mathcal{D}_1$ such that (59) holds.

(ii)

$$c_1(x_1) = c_2(x_2) \quad (60)$$

Here $T_{(x_1, x_2)}\mathcal{R} \subset T_{x_1}\mathcal{X}_1 \times T_{x_2}\mathcal{X}_2$ denotes the tangent space to \mathcal{R} at the point $(x_1, x_2) \in \mathcal{R}$.

Furthermore, using the theory of nonlinear controlled invariance, in particular the theory of controlled invariant submanifolds, see e.g. [12, 17], we derive the following extension of Theorem 2.10.

Theorem 7.2. *A submanifold $\mathcal{R} \subset \mathcal{X}_1 \times \mathcal{X}_2$ is a bisimulation relation between Σ_1 and Σ_2 if and only if for all $(x_1, x_2) \in \mathcal{R}$*

$$\begin{aligned}
(a) \quad & T_{(x_1, x_2)}\mathcal{R} + \text{im} \begin{bmatrix} g_1(x_1) \\ 0 \end{bmatrix} = T_{(x_1, x_2)}\mathcal{R} + \text{im} \begin{bmatrix} 0 \\ g_2(x_2) \end{bmatrix} =: \mathcal{R}_e(x_1, x_2) \\
(b) \quad & \begin{bmatrix} a_1(x_1) \\ a_2(x_2) \end{bmatrix} \in \mathcal{R}_e(x_1, x_2) \\
(c) \quad & \text{im} \begin{bmatrix} b_1(x_1) \\ b_2(x_2) \end{bmatrix} \subset \mathcal{R}_e(x_1, x_2) \\
(d) \quad & c_1(x_1) = c_2(x_2)
\end{aligned} \tag{61}$$

Remark 7.3. Note that $\mathcal{R}_e(x_1, x_2)$ is a linear subspace of the tangent space $T_{x_1}\mathcal{X}_1 \times T_{x_2}\mathcal{X}_2$, which is defined for all $(x_1, x_2) \in \mathcal{R}$.

Remark 7.4. For simplicity of presentation we have restricted ourselves to nonlinear systems (58), instead of the more general form $\dot{x}_i = f(x_i, u_i, d_i)$, $y_i = h(x_i, u_i)$. Proposition 7.1 easily generalizes by considering instead of (59) the conditions $(f_1(x_1, u, d_1), f_2(x_2, u, d_2)) \in T_{(x_1, x_2)}\mathcal{R}$ and $h_1(x_1, u) = h_2(x_2, u)$. An appropriate generalization of Theorem 7.2 can be obtained along the lines of [17] (Chapter 13).

For computing the maximal nonlinear bisimulation relation one may use the following version of the algorithm to compute the maximal controlled invariant submanifold, cf. [17].

Algorithm 7.5. *Given two nonlinear dynamical systems Σ_1 and Σ_2 as in (58). Define the following sequence $\mathcal{R}^j, j = 0, 1, 2, \dots$, of subsets of $\mathcal{X}_1 \times \mathcal{X}_2$, and assume that they are all submanifolds (possibly empty)*

$$\begin{aligned}
\mathcal{R}^0 &= \mathcal{X}_1 \times \mathcal{X}_2 \\
\mathcal{R}^1 &= \{(x_1, x_2) \in \mathcal{R}^0 \mid c_1(x_1) = c_2(x_2)\} \\
\mathcal{R}^2 &= \{(x_1, x_2) \in \mathcal{R}^1 \mid \begin{bmatrix} a_1(x_1) \\ a_2(x_2) \end{bmatrix} + \text{im} \begin{bmatrix} g_1(x_1) \\ 0 \end{bmatrix} \subset T_{(x_1, x_2)}\mathcal{R}^1 + \text{im} \begin{bmatrix} 0 \\ g_2(x_2) \end{bmatrix}, \\
&\quad \left. \begin{bmatrix} a_1(x_1) \\ a_2(x_2) \end{bmatrix} + \text{im} \begin{bmatrix} 0 \\ g_2(x_2) \end{bmatrix} \subset T_{(x_1, x_2)}\mathcal{R}^1 + \text{im} \begin{bmatrix} g_1(x_1) \\ 0 \end{bmatrix} \right\} \\
&\vdots \\
\mathcal{R}^{j+1} &= \{(x_1, x_2) \in \mathcal{R}^j \mid \begin{bmatrix} a_1(x_1) \\ a_2(x_2) \end{bmatrix} + \text{im} \begin{bmatrix} g_1(x_1) \\ 0 \end{bmatrix} \subset T_{(x_1, x_2)}\mathcal{R}^j + \text{im} \begin{bmatrix} 0 \\ g_2(x_2) \end{bmatrix}, \\
&\quad \left. \begin{bmatrix} a_1(x_1) \\ a_2(x_2) \end{bmatrix} + \text{im} \begin{bmatrix} 0 \\ g_2(x_2) \end{bmatrix} \subset T_{(x_1, x_2)}\mathcal{R}^j + \text{im} \begin{bmatrix} g_1(x_1) \\ 0 \end{bmatrix} \right\}
\end{aligned} \tag{62}$$

Remark 7.6. For a discussion of the condition that \mathcal{R}^j at every step of the algorithm is a submanifold we refer to [17] (Chapter 11). For the non-smooth case a similar type of algorithm

has been provided in the context of viability theory; see e.g. [3] and the references quoted in there.

Proposition 7.7. 1. $\mathcal{R}^0 \supset \mathcal{R}^1 \supset \mathcal{R}^2 \supset \dots \supset \mathcal{R}^j \supset \mathcal{R}^{j+1} \supset \dots$

2. There exists a finite k such that $\mathcal{R}^k = \mathcal{R}^{k+1} =: \mathcal{R}^*$, and then $\mathcal{R}^j = \mathcal{R}^*$ for all $j \geq k$.

3. \mathcal{R}^* is the maximal submanifold of $\mathcal{X}_1 \times \mathcal{X}_2$ (possibly empty) satisfying properties (61a,b,d).

Corollary 7.8. Suppose that Algorithm 7.5 is well-defined, that is, the subsets \mathcal{R}^j are all submanifolds. Then Σ_1 and Σ_2 are bisimilar if and only if \mathcal{R}^* satisfies (61c) and $\pi_i(\mathcal{R}^*) = \mathcal{X}_i, i = 1, 2$.

Reduction of a nonlinear system

$$\begin{aligned} \dot{x} &= a(x) + b(x)u + g(x)d, & x \in \mathcal{X}, & u \in \mathcal{U}, & d \in \mathcal{D} \\ \Sigma : & & & & \\ y &= c(x) & & & y \in \mathcal{Y} \end{aligned} \tag{63}$$

to a nonlinear system with lower-dimensional state space $\mathcal{X}_{\mathcal{R}}$ can be achieved, similarly to the linear theory exposed in Section 4, by considering a nonlinear bisimulation relation \mathcal{R} between Σ and itself. Such a bisimulation relation is given by a submanifold $\mathcal{R} \subset \mathcal{X} \times \mathcal{X}$ satisfying for all $(x_1, x_2) \in \mathcal{R}$

$$\begin{aligned} (a) \quad & T_{(x_1, x_2)}\mathcal{R} + \text{im} \begin{bmatrix} g(x_1) \\ 0 \end{bmatrix} = T_{(x_1, x_2)}\mathcal{R} + \text{im} \begin{bmatrix} 0 \\ g(x_2) \end{bmatrix} =: \mathcal{R}_e(x_1, x_2) \\ (b) \quad & \begin{bmatrix} a(x_1) \\ a(x_2) \end{bmatrix} \in \mathcal{R}_e(x_1, x_2) \\ (c) \quad & \text{im} \begin{bmatrix} b(x_1) \\ b(x_2) \end{bmatrix} \subset \mathcal{R}_e(x_1, x_2) \\ (d) \quad & c(x_1) = c(x_2) \end{aligned} \tag{64}$$

Similarly to the linear theory in Section 4 we assume that \mathcal{R} defines an *equivalence* relation on \mathcal{X} . Then for all $x \in \mathcal{X}$ we may define the equivalence classes

$$\begin{aligned} \mathcal{F}_{\mathcal{R}}^x &= \{x' \in \mathcal{X} \mid (x', x) \in \mathcal{R}\} \\ &= \pi(\mathcal{R} \cap \{(x', x) \mid x' \in \mathcal{X}\}) \end{aligned} \tag{65}$$

where π denotes the projection on the first component. Throughout we assume that the subsets $\mathcal{F}_{\mathcal{R}}^x$ of \mathcal{X} are *submanifolds* of \mathcal{X} for every $x \in \mathcal{X}$. (Under generic transversality conditions this will be the case since $\mathcal{F}_{\mathcal{R}}^x$ is defined as the intersection of submanifolds.)

Since \mathcal{R} is assumed to be an equivalence relation it follows that $\mathcal{F}_{\mathcal{R}}^x, x \in \mathcal{X}$, defines a *foliation* $F_{\mathcal{R}}$ of \mathcal{X} in the following sense:

$$\begin{aligned} (i) \quad & x \in \mathcal{F}_{\mathcal{R}}^x \\ (ii) \quad & y \in \mathcal{F}_{\mathcal{R}}^x \Leftrightarrow x \in \mathcal{F}_{\mathcal{R}}^y \\ (iii) \quad & \mathcal{F}_{\mathcal{R}}^x \cap \mathcal{F}_{\mathcal{R}}^y \neq \emptyset \Rightarrow \mathcal{F}_{\mathcal{R}}^x = \mathcal{F}_{\mathcal{R}}^y \end{aligned} \tag{66}$$

Now let us consider conditions (64) in more detail. First of all, condition (64b) implies that there exists a vector $\alpha(x_1, x_2)$ such that

$$\begin{bmatrix} a'(x_1) \\ a'(x_2) \end{bmatrix} := \begin{bmatrix} a(x_1) \\ a(x_2) \end{bmatrix} + \begin{bmatrix} g(x_1) \\ 0 \end{bmatrix} \alpha(x_1, x_2) \in T_{(x_1, x_2)}\mathcal{R} \quad (67)$$

Hence the vector field $\begin{bmatrix} a'(x_1) \\ a'(x_2) \end{bmatrix}$ is tangent to \mathcal{R} for every $(x_1, x_2) \in \mathcal{R}$. This means that the *flow* of this vector field when starting at $(x_1, x_2) \in \mathcal{R}$ will *remain* in \mathcal{R} for all $t > 0$ for which it is defined. This implies that the flow of this vector field when projected to the first factor of $\mathcal{X} \times \mathcal{X}$ leaves the foliation $\mathcal{F}_{\mathcal{R}}$ invariant. Using the theory of invariant distributions (see e.g. [12, 17]) this is known to be equivalent to the condition

$$[a', D_{\mathcal{R}}] \subset D_{\mathcal{R}} \quad (68)$$

where $D_{\mathcal{R}}$ is the integrable distribution on \mathcal{X} corresponding to $\mathcal{F}_{\mathcal{R}}$, i.e.,

$$D_{\mathcal{R}}(x) := T_x \mathcal{F}_{\mathcal{R}}^x, \quad x \in \mathcal{X} \quad (69)$$

Similar statements hold with respect to invariance of $D_{\mathcal{R}}$ with respect to the columns of b and g . Using the theory of controlled invariant distributions [12, 17] one then arrives at the following nonlinear analogue of Theorem 4.2.

Proposition 7.9. *Let the submanifold $\mathcal{R} \subset \mathcal{X} \times \mathcal{X}$ define an equivalence relation on \mathcal{X} . Define the integrable distribution $D_{\mathcal{R}}$ on \mathcal{X} by (69). Then \mathcal{R} is a bisimulation relation between Σ and itself if and only if $D_{\mathcal{R}}$ satisfies*

$$\begin{aligned} [a, D_{\mathcal{R}}] &\subset D_{\mathcal{R}} + \text{im } g \\ [b, D_{\mathcal{R}}] &\subset D_{\mathcal{R}} + \text{im } g \\ [g, D_{\mathcal{R}}] &\subset D_{\mathcal{R}} + \text{im } g \\ D_{\mathcal{R}} &\subset \ker dc \end{aligned} \quad (70)$$

Conditions (70) express that the distribution $D_{\mathcal{R}}$ is controlled invariant (for a and the columns of b with respect to the ‘input’ distribution $\text{im } g(x)$). It follows that the foliation $\mathcal{F}_{\mathcal{R}}$ (or equivalently the distribution $D_{\mathcal{R}}$) can be factored out to obtain a reduced nonlinear system $\Sigma_{\mathcal{R}}$ (assuming that the quotient space is again a manifold, see for a discussion of this e.g. [8]). We leave the details to the reader.

Example 7.10. Consider the nonlinear system (cf. [28])

$$\begin{aligned} \dot{x}^1 &= x^2 d^1 \\ \Sigma : \dot{x}^2 &= d^2 \\ y &= x^1 \end{aligned} \quad (71)$$

This system is *not* reducible by bisimulation to the system

$$\begin{aligned} \dot{z} &= d \\ y &= z \end{aligned} \quad (72)$$

(although their C^∞ or piecewise C^∞ external behaviors are equal!). Indeed, the distribution $D_{\mathcal{R}} := \text{span} \left\{ \frac{\partial}{\partial x^2} \right\}$ does not satisfy (70). Reason is that Σ given by (71) has a ‘deadlock’ behavior for $x^2 = 0$, which is not present in the system given by (72). Note that if we replace the first equation of (71) by $\dot{x}^1 = x^2 + d^1$ then Σ is (linearly) bisimilar to (72).

Finally we obtain the following nonlinear (partial) analogue of Theorem 6.3 for the case that $g_i = 0$ (the ‘deterministic’ case).

Proposition 7.11. *A submanifold $\mathcal{R} \subset \mathcal{X}_1 \times \mathcal{X}_2$ is a bisimulation relation between Σ_1 and Σ_2 for $g_i = 0$ if and only if for all $(x_1, x_2) \in \mathcal{R}$*

$$\begin{aligned}
 (a) \quad & \begin{bmatrix} a_1(x_1) \\ a_2(x_2) \end{bmatrix} \in T_{(x_1, x_2)} \mathcal{R} \\
 (b) \quad & \text{im} \begin{bmatrix} b_1(x_1) \\ b_2(x_2) \end{bmatrix} \subset T_{(x_1, x_2)} \mathcal{R} \\
 (c) \quad & c_1(x_1) = c_2(x_2)
 \end{aligned} \tag{73}$$

Furthermore, \mathcal{R} is a bisimulation relation if and only if for all $(x_1, x_2) \in \mathcal{R}$

$$H_1(x_1) = H_2(x_2) \tag{74}$$

for all corresponding functions $H_i := L_{X_i^1} L_{X_i^2} \cdots L_{X_i^k} c_i$, where X_i^j are the vector fields given by a_i or the columns of b_i , in the observation space [17] of $\Sigma_i, i = 1, 2$. Finally, the maximal bisimulation relation between Σ_1 and Σ_2 equals the set of points (x_1, x_2) satisfying (74).

Proof. Obviously, (73) is the specialization of (64) to the present case. By invariance of the observation space of Σ_i under a_i and the columns of b_i it follows that satisfaction of (74) implies (73). ■

8 Conclusions and outlook

In this paper we have defined and studied a notion of bisimulation for continuous dynamical systems, motivated by the theory of bisimulation for concurrent processes and by previously obtained results by Pappas and co-authors. Furthermore we have made explicit the relations with well-known system-theoretic notions of state space equivalence and reduction. The notion of bisimulation appears to be a notion which *unifies* the concepts of *state space equivalence* and *state space reduction*, and which allows to study equivalence of systems with *non-minimal state space dimension*. Classical results concerning reduction and state space equivalence for minimal deterministic systems are readily and elegantly recovered. A crucial tool in all this is the notion of controlled invariance as developed in linear and nonlinear geometric control theory.

Compared with classical systems theory a new twist to the problem is given by the idea of considering *non-deterministic* continuous dynamical systems. For concurrent discrete processes the advantages of allowing non-determinism are clear [15, 11], but for continuous dynamical systems such arguments have not been explicitated except in the context of abstraction. Indeed, in the area of verification of hybrid systems it is natural to look for notions of abstraction which allow to extend methods for verification of discrete processes to the hybrid case, see [10, 2, 13, 14, 30]. This naturally leads to considering ways of abstracting

continuous dynamical systems. Apart from abstraction we believe that there are other compelling reasons to study some type of ‘non-determinism’ in continuous dynamical systems. In particular, it would be interesting to investigate if uncertainty and robustness issues can be fruitfully cast in this framework. Another line of research could be to replace the generator for non-determinism d by a stochastic process (although this would presumably also alter the definition of equivalence). Finally, we do not necessarily have to think about d as disturbances; instead d may be the interconnection variables when the system is interconnected to another system, in particular to a controller system. Current research is aimed at exploring such ideas in the extension of the results on achievable closed-loop behavior obtained in [23, 26] to the situation where the closed-loop system is specified up to bisimulation. Notice in this respect that bisimulation for continuous dynamical systems satisfies the basic rule of modularity: if Σ_1 is bisimilar to Σ_2 then also the interconnection of Σ_1 with any controller system will be bisimilar to the interconnection of Σ_2 with the same controller system.

Apart from the interest per se in the equivalence concept for dynamical systems studied in this paper a major aim of this work is eventually to come up with a general notion of bisimulation for (classes of) *hybrid systems*, that is, open systems with interacting continuous and discrete dynamics. As indicated above, already a substantial amount of work has been done in this area. One way of approaching this problem (especially if one is looking for finite or almost finite bisimulations of hybrid systems) is to abstract away from the time, see e.g. [10, 2, 19, 9]. On the other hand, another approach would be to *combine* the notion of bisimulation as explored in the current paper with the established notion of bisimulation for concurrent processes into a notion of *structural bisimulation* for hybrid systems. Initial steps in this direction have been taken in [24, 25].

Finally, a generalization to the bisimulation of continuous dynamical systems as exposed in the present paper is to suppress the a priori distinction of the external variables into inputs u and outputs y . This can be achieved by considering a behavioral version of (1), e.g. given in pencil representation as

$$\begin{aligned} E\dot{z} &= Az \\ w &= Hz \end{aligned} \tag{75}$$

where now w denotes the *whole* vector of external variables (not split into u and y). Note that (1) may be put into the form (75) by first pre-multiplying the first line of (1) by an annihilating matrix G^\perp ($G^\perp G = 0$) of maximal rank, to obtain $G^\perp \dot{x} = G^\perp Ax + G^\perp Bu$, and then considering the extended state vector $z = (x, u)$. Generalization of the bisimulation theory to this class of systems is pursued in [25]. An additional and important advantage of (75) is that it may also include algebraic constraints on the state variables, and thus is eminently suited to model systems as resulting from interconnection of systems, or multimodal systems arising from varying (event-driven) state constraints [27].

Acknowledgements

Initial discussions with George Pappas and Paulo Tabuada (University of Pennsylvania) and ongoing conversations with Jan Willem Polderman, Rom Langerak, Agung Julius and Stefan Strubbe (all University of Twente) are highly appreciated.

References

- [1] R. Abraham, J.E. Marsden, T. Ratiu, *Manifolds, Tensor Analysis, and Applications*, Applied Mathematical Sciences 75, 2nd edition, Springer, New York, 1988.
- [2] R. Alur, T.A. Henzinger, G. Lafferriere, G.J. Pappas, Discrete abstractions of hybrid systems, *Proceedings of the IEEE*, vol. 88, pp. 971–984, 2000.
- [3] J.-P. Aubin, J. Lygeros, M. Quincampoix, Shankar Sastry, N. Seube, Impulse differential inclusions: A viability approach to hybrid systems, *IEEE Transactions on Automatic Control*, 47(1), pp. 2–20, 2002.
- [4] G. Basile, G. Marro, *Controlled and conditioned invariants in linear system theory*, Prentice Hall, Englewood Cliffs, 1992.
- [5] R.W. Brockett, *Finite Dimensional Linear Systems*, John Wiley, New York, 1970.
- [6] F.M. Callier, C.A. Desoer, *Linear System Theory*, Springer-Verlag, New York, 1991.
- [7] E.M. Clarke, Jr., O. Grumberg, D.A. Peled, *Model Checking*, The MIT Press, Cambridge, Massachusetts, 2002.
- [8] V.I. Elkin, *Reduction of nonlinear control systems : a differential geometric approach*, Kluwer, Dordrecht, 1999.
- [9] E. Haghverdi, P. Tabuada, G.J. Pappas, Unifying bisimulation relations for discrete and continuous systems, *Category Theory and Computer Science, Electronic Notes in Theoretical Computer Science*, August 2002.
- [10] T.A. Henzinger, “Hybrid automata with finite bisimulations”, in *ICALP 95: Automata, Languages, and programming*, Eds. Z. Fulop, F. Gecseg, Lecture Notes in Computer Science 944, pp. 324–335, Springer, New York, 1995
- [11] H. Hermanns, *Interactive Markov Chains*, Lecture Notes in Computer Science 2428, Springer, 2002.
- [12] A. Isidori, *Nonlinear Control Systems*, Third edition, Springer, London, 1995.
- [13] G. Lafferriere, G.J. Pappas, S. Sastry, “Hybrid systems with finite bisimulations”, in *Hybrid Systems V*, eds. P. Antsaklis, W. Kohn, M. Lemmon, A. Nerode, S. Sastry, Lecture Notes in Computer Science, Springer, 1998.
- [14] G. Lafferriere, G.J. Pappas, S. Sastry, “O-minimal hybrid systems”, *Math. Contr. Signals, Syst.*, vol. 13, pp. 1-21, 2000.
- [15] R. Milner, *Communication and Concurrency*, Prentice Hall International Series in Computer Science, 1989.
- [16] R. Milner, *Communication and Mobile Systems: the π - Calculus*, Cambridge University Press, Cambridge, UK, 1999.
- [17] H. Nijmeijer, A.J. van der Schaft, *Nonlinear Dynamical Control Systems*, Springer-Verlag, New York, 1990 (4th printing 1998), p. xiii+467.

- [18] G.J. Pappas, G. Lafferriere, S. Sastry, Hierarchically consistent control systems, *IEEE Transactions on Automatic Control*, 45(6), pp. 1144–1160, 2000.
- [19] G.J. Pappas, Bisimilar linear systems, *Automatica*, 39(12), pp. 2035–2047, 2003.
- [20] G.J. Pappas, S. Simic, Consistent abstractions of affine control systems, *IEEE Transactions on Automatic Control*, 47, pp. 745–756, 2002.
- [21] D. Park, Concurrency and automata on infinite sequences. In P. Deussen, Editor, *Fifth GI Conference on Theoretical Computer Science*, vol. 104 of *Lecture Notes in Computer Science*, Springer, 1981.
- [22] J.W. Polderman, J.C. Willems, *Introduction to Mathematical Systems Theory*, Springer-Verlag, New York, 1998.
- [23] A.J. van der Schaft, Achievable behavior of general systems. *Systems & Control Letters*, 49, pp. 141–149, 2003.
- [24] A.J. van der Schaft, Bisimulation of dynamical systems, in *Hybrid Systems: Computation and Control*, eds. R. Alur, G.J. Pappas, Lecture Notes in Computer Science 2293, pp. 555–569, Springer, 2004.
- [25] A.J. van der Schaft, “Equivalence of hybrid dynamical systems”, *Proceedings of the Sixteenth International Symposium on Mathematical Theory of Networks and Systems (MTNS2004)*, Leuven, Belgium, July 5–9, 2004.
- [26] A.J. van der Schaft, A.A. Julius, “Achievable behavior by composition”, *Proc. 41st IEEE Conf. Decision and Control*, Las Vegas, Nevada, December 2002.
- [27] A.J. van der Schaft, J.M. Schumacher, *An Introduction to Hybrid Dynamical Systems*, Springer Lecture Notes in Control and Information Sciences, Vol.251, Springer-Verlag, London, 2000.
- [28] A.J. van der Schaft, J.M. Schumacher, “Compositionality issues in discrete, continuous, and hybrid systems”, *Int. J. Robust and Nonlinear Control*, vol. 11, pp. 417–434, 2001.
- [29] P. Tabuada, G.J. Pappas, “Bisimilar control affine systems”, *Systems and Control Letters*, 52, pp. 49–58, 2004.
- [30] P. Tabuada, G.J. Pappas, P. Lima, “Composing abstractions of hybrid systems”, in *Hybrid Systems: Computation and Control*, eds. C. Tomlin, M.R. Greenstreet, Lecture Notes in Computer Science, pp. 436–450, Springer, 2002.
- [31] W.M. Wonham, *Linear multivariable control: a geometric approach*, Springer, New York, Third Edition, 1985.